

Lesson Summary

Some proportions are linear equations in disguise and are solved the same way we normally solve proportions.

When multiplying a fraction with more than one term in the numerator and/or denominator by a number, put the expressions with more than one term in parentheses so that you remember to use the distributive property when transforming the equation. For example:

$$\frac{x+4}{2x-5} = \frac{3}{5}$$

$$5(x+4) = 3(2x-5).$$

The equation $5(x+4) = 3(2x-5)$ is now clearly a linear equation and can be solved using the properties of equality.

Problem Set

Solve the following equations of rational expressions, if possible. If an equation cannot be solved, explain why.

1. $\frac{5}{6x-2} = \frac{-1}{x+1}$

6. $\frac{2x+5}{2} = \frac{3x-2}{6}$

2. $\frac{4-x}{8} = \frac{7x-1}{3}$

7. $\frac{6x+1}{3} = \frac{9-x}{7}$

3. $\frac{3x}{x+2} = \frac{5}{9}$

8. $\frac{\frac{1}{3}x-8}{12} = \frac{-2-x}{15}$

4. $\frac{\frac{1}{2}x+6}{3} = \frac{x-3}{2}$

9. $\frac{3-x}{1-x} = \frac{3}{2}$

5. $\frac{7-2x}{6} = \frac{x-5}{1}$

10. In the diagram below, $\triangle ABC \sim \triangle A'B'C'$. Determine the lengths of \overline{AC} and \overline{BC} .

